

Joint numerical ranges, quantum maps, and joint numerical shadows

Eugene Gutkin^{a,b}, Karol Życzkowski^{c,d}

^a*Department of Mathematics, Nicolaus Copernicus University, Chopina 12/18, 87-100 Toruń, Poland*

^b*Institute of Mathematics, Polish Academy of Sciences, Śniadeckich 8, 00-956 Warszawa 10, Poland*

^c*Instytut Fizyki im. Smoluchowskiego, Uniwersytet Jagielloński, Reymonta 4, 30-059 Kraków, Poland*

^d*Centrum Fizyki Teoretycznej, Polska Akademia Nauk, Aleja Lotników 32/44, 02-668 Warszawa, Poland*

Abstract

We associate with k hermitian $N \times N$ matrices a probability measure on $\mathbb{R}^k$. It is supported on the joint numerical range of the k -tuple of matrices. We call this measure the joint numerical shadow of these matrices. Let $k = 2$. A pair of hermitian $N \times N$ matrices defines a complex $N \times N$ matrix. The joint numerical range and the joint numerical shadow of the pair of hermitian matrices coincide with the numerical range and the numerical shadow, respectively, of this complex matrix. We study relationships between the dynamics of quantum maps on the set of quantum states, on one hand, and the numerical ranges, on the other hand. In particular, we show that under the identity resolution assumption on Kraus operators defining the quantum map, the dynamics shrinks numerical ranges.

Keywords: joint numerical range, positive matrices, quantum maps, quantum states
2000 MSC: 47A12, 81P16, 51M15

1. Introduction

We denote by $\mathcal{H}_N$ the space $\mathbb{C}^N$ endowed with the standard scalar product $\langle \varphi | \psi \rangle$. Thus, $\mathcal{H}_N$ is the Hilbert space of N dimensions. Let A be an operator on $\mathcal{H}_N$. Its *numerical range* (also called the *field of values*) $W(A) \subset \mathbb{C}$ is the set of numbers $z = \langle \psi | A | \psi \rangle$, where $\langle \psi | \psi \rangle = 1$ [1, 2]. A crucial fact, conjectured by O. Toeplitz in 1918 [3] and proved by F. Hausdorff in 1919 [4] is that $W(A)$ is convex. See [5] for a modern exposition and historical remarks. If A is a normal operator, then $W(A)$ is the convex hull of the spectrum of A [1], thus a convex polygon. For generic A the boundary $\partial W(A)$ is smooth [6]. For $N = 2$, $\partial W(A)$ is a (possibly degenerate) ellipse [3, 7]. See [8] for a classification of numerical ranges when $N = 3$.

In recent years, numerical range and its generalizations found novel applications in quantum mechanics, in particular, in the theory of quantum information [9, 10, 11]. We will freely use the physics terminology in what follows. *Pure quantum states* are the hermitian projections of $\mathcal{H}_N$ onto one-dimensional subspaces $\mathbb{C}|\psi\rangle$. The set Ω_N of pure quantum states is naturally isomorphic to the complex projective space $\mathbb{C}P^{N-1}$. A *density matrix* is a positive operator $\rho \geq 0$ on $\mathcal{H}_N$ satisfying $\text{Tr} \rho = 1$. A pure quantum

Preprint submitted to Elsevier

July 6, 2012

state is a density matrix, and any density matrix is a convex combination of pure quantum states. In the physics terminology, density matrices are the *quantum states*. The set $\mathcal{Q}_N$ of quantum states is convex, and $\Omega_N \subset \mathcal{Q}_N$ is the set of its extremal points. The elements of $\mathcal{Q}_N \setminus \Omega_N$ are *mixed quantum states*.

Applications of the numerical range to quantum information are based on the interpretation of $W(A)$ as a plane projection of Ω_N [12]. For instance, in the *one qubit case*, $N = 2$, we have $\Omega_2 = \mathbf{CP}^1 = S^2$ (the *Bloch sphere*), and $\mathcal{Q}_2 = B^3 \subset \mathbf{R}^3$ (the *Bloch ball*). A projection of the two-sphere is a possibly degenerate ellipse. This simple observation is the base for the well known facts about numerical ranges of 2×2 matrices [7].

Let A be an operator on $\mathcal{H}_N$. Its *numerical shadow* $P_A(z)$ is a probability distribution on $\mathbb{C}$ supported on $W(A)$ [15, 13, 14]. Let $|\psi\rangle$ be distributed on the unit sphere in $\mathcal{H}_N$ according to the unique rotationally invariant probability measure (i.e., the Haar measure). Then $P_A(z)$ is the probability density of $z = \langle \psi | A | \psi \rangle$. Let $N = 2$. Then $P_A(z)$ is the density of the plane shadow of the sphere under a light beam. This explains the term “numerical shadow” [15].

Let $A_1, \dots, A_m$ be hermitian operators on $\mathcal{H}_N$. Their *joint numerical range* (JNR) $W(A_1, \dots, A_m) \subset \mathbb{R}^m$ [17] is defined as follows:

$$W(A_1, \dots, A_m) = \{(\langle \psi | A_1 | \psi \rangle, \dots, \langle \psi | A_m | \psi \rangle) : \langle \psi | \psi \rangle = 1\}. \quad (1)$$

Let $m = 2$. Then $W(A_1, A_2) = W(A_1 + iA_2)$. Thus, JNR generalizes the notion of the numerical range of a complex operator. If $m > 2$, then $W(A_1, \dots, A_m) \subset \mathbb{R}^m$ is not necessarily convex [17]. For instance, JNR of the triple of Pauli matrices is the Bloch sphere. We will elaborate on this observation in section 2.

In this work we establish and explore a relationship between the notion of the joint numerical range and the set of quantum states. In Theorem 1 we show that the JNR of an m -tuple of hermitian operators is a suitable linear projection of the set of pure quantum states. Using this observation, we associate with the joint numerical range a probability measure on $\mathbf{R}^m$, the *joint numerical shadow* of an m -tuple of hermitian operators. See section 4. This extends the concept of numerical shadow to arbitrary dimensions [15, 13, 14, 16]. Let Φ denote the discrete dynamics on the set of quantum states defined by a set of k Kraus operators satisfying the identity resolution equation (11). We analyze the effect of the dynamics Φ on the numerical range. In Corollary 1 we obtain an inclusion of the numerical range of $\Phi(A)$ into the numerical range of A .

2. Joint numerical ranges

Let $\mathcal{L}_N$ (resp. $\mathcal{M}_N$, $\mathcal{P}_N$, $\mathcal{Q}_N$) denote the space of all (resp. hermitian, positive definite, positive definite with trace 1) linear operators on $\mathcal{H}_N$. Let $\Omega_N \subset \mathcal{Q}_N$ be the set of rank one projections. Then $\mathcal{L}_N$ endowed with the scalar product

$$(A, B) = \text{tr}(AB^*) \quad (2)$$

is a Hilbert space of N^2 dimensions, and $\mathcal{M}_N \subset \mathcal{L}_N$ is a real subspace of N^2 dimensions on which the scalar product (2) induces the structure of a euclidean space. The set

$\mathcal{P}_N \subset \mathcal{M}_N$ is a closed convex cone. Its interior consists of strictly positive operators, $\rho > 0$, and its apex is the zero operator. The set $\mathcal{Q}_N$ is the intersection of $\mathcal{P}_N$ and the hyperplane $\{\text{Tr}\rho = 1\}$. It is a bounded convex region (i.e., has nonempty interior) in the $(N^2 - 1)$ -dimensional affine space, and $\Omega_N \subset \mathcal{Q}_N$ is the set of its extremal points. For a unit vector $\psi \in \mathcal{H}_N$ we set $\rho_\psi = |\psi\rangle\langle\psi| \in \Omega_N$. The manifold $\mathbb{C}P^{N-1}$ is the quotient of the unit sphere $S(\mathcal{H}_N)$ by the natural linear action of the unit circle $S^1 \subset \mathbb{C}$. The map $\psi \mapsto \rho_\psi$ yields an isomorphism of $\mathbb{C}P^{N-1}$ and Ω_N . In what follows we will identify $\mathbb{C}P^{N-1}$ and Ω_N via this isomorphism.

Let $A_1, \dots, A_m \in \mathcal{M}_N$ be arbitrary. The mapping from $\mathbb{C}P^{N-1}$ to $\mathbb{R}^m$ given by

$$\text{jnr}_{(A_1, \dots, A_m)} : |\psi\rangle \mapsto (\langle\psi|A_1|\psi\rangle, \dots, \langle\psi|A_m|\psi\rangle). \quad (3)$$

is the *joint numerical range map*. The range $W(A_1, \dots, A_m)$ of this map is the *joint numerical range* (JNR) of operators $A_1, \dots, A_m$. By the isomorphism $\mathbb{C}P^{N-1} \simeq \Omega_N$, we have $\text{jnr}_{(A_1, \dots, A_m)} : \Omega_N \rightarrow \mathbb{R}^m$.

We will recall a few basic notions in affine geometry. By a vector space we will mean a finite dimensional real vector space. Let V be a vector space. A set $H \subset V$ is an *affine subspace* if there is a linear subspace $H_0 \subset V$ and a vector $h_0 \in H$ such that $H = H_0 + h_0$. Let $G \subset U, H \subset V$ be affine subspaces. Let $G_0 \subset U, H_0 \subset V$ be the corresponding linear subspaces. A map $Af : G \rightarrow H$ is an *affine isomorphism* if there is a linear isomorphism $Af_0 : G_0 \rightarrow H_0$ and vectors $h_0 \in H, g_0 \in G$ so that

$$Af(g) = Af_0(g - g_0) + h_0.$$

Definition 1. Let U, V be vector spaces, and let $X \subset U, Y \subset V$ be arbitrary sets. They are *affinely isomorphic* if there exist affine subspaces $H \subset U, G \subset V$ such that $X \subset G, Y \subset H$, and an affine isomorphism $A : G \rightarrow H$ such that $A(X) = Y$. The induced map $A : X \rightarrow Y$ is an *affine isomorphism of X onto Y* .

Note that linear isomorphism of sets are the special cases in the above setting when the subspaces and the maps in question are, actually, linear. We will not distinguish between the linear and affine situations in what follows.

We will now expose a general topic in linear algebra. Let U, V be vector spaces. We assume that U is a euclidean space with the scalar product $(\cdot, \cdot)$. Let $m \geq 1$. With any nonzero vectors $u_1, \dots, u_m \in U$ and $v_1, \dots, v_m \in V$ we associate a linear operator $L : U \rightarrow V$ by

$$L(u) = \sum_{i=1}^m (u, u_i) v_i. \quad (4)$$

Let $\mathcal{A} \subset U$ (resp. $\mathcal{B} \subset V$) be the subspace spanned by $u_1, \dots, u_m$ (resp. $v_1, \dots, v_m$). We will need a simple lemma about the operator in equation (4). The following claim should be in the literature. We do not give a reference, but the proof is straightforward, and we leave it to the reader.

Lemma 1. 1. Let $\dim \mathcal{A} = p \leq m$. Assume without loss of generality that the vectors $u_1, \dots, u_p$ span $\mathcal{A}$. Then there are vectors $\tilde{v}_1, \dots, \tilde{v}_p \in \mathcal{B}$ such that the linear operator in equation (4) satisfies

$$L(u) = \sum_{i=1}^p (u, u_i) \tilde{v}_i. \quad (5)$$

2. Let $\dim \mathcal{B} = q \leq m$. Assume without loss of generality that the vectors $v_1, \dots, v_q$ span $\mathcal{B}$. Then there are vectors $\tilde{u}_1, \dots, \tilde{u}_q \in \mathcal{A}$ such that the linear operator in equation (4) satisfies

$$L(u) = \sum_{i=1}^q (u, \tilde{u}_i) v_i. \quad (6)$$

3. If $p = m$ (resp. $q = m$) then $\tilde{v}_i = v_i$ (resp. $\tilde{u}_i = u_i$) for $1 \leq i \leq m$.

The proposition below is immediate from Lemma 1.

Proposition 1. Let the setting be as in Lemma 1, and let $Pr_{\mathcal{A}} : U \rightarrow \mathcal{A}$ be the orthogonal projection. Then i) There is a unique linear operator $M : \mathcal{A} \rightarrow \mathcal{B}$ such that $L = M \circ Pr_{\mathcal{A}}$; ii) If $\dim \mathcal{A} = m$ (resp. $\dim \mathcal{B} = m$) then M is surjective (resp. injective); iii) If $\dim \mathcal{A} = \dim \mathcal{B} = m$ then M is an isomorphism.

Let $G, H \subset U$ be affine subspaces in a vector space. We say that they are *parallel* if $H = G + u_0$ for some $u_0 \in U$.

Corollary 1. Let U, V be vector spaces, and let $L : U \rightarrow V$ be as in equation (4). Let $\mathcal{A} \subset U, \mathcal{B} \subset V$ be the subspaces spanned by the vectors $u_1, \dots, u_m$ and $v_1, \dots, v_m$ respectively. Let $M : \mathcal{A} \rightarrow \mathcal{B}$ be the operator from Proposition 1.

Let $G \subset U$ be an affine subspace containing a subspace parallel to $\mathcal{A}$. Let $\Gamma \subset G$ be an arbitrary set. If M is injective, then $Pr_{\mathcal{A}}(\Gamma)$ and $L(\Gamma)$ are affinely isomorphic. **Proof.** Let $u_0 \in U$ be such that $\mathcal{A} = G + u_0$. Let $\mathcal{B}_0 \subset \mathcal{B}$ be the range of M . Thus, $M : \mathcal{A} \rightarrow \mathcal{B}_0$ is a linear isomorphism. By Proposition 1, we have

$$L(\Gamma + u_0) = M Pr_{\mathcal{A}}(\Gamma + u_0) = M(\Gamma + u_0) \subset \mathcal{B}_0.$$

The mapping $u \mapsto L(u + u_0)$ induces an affine isomorphism of G and $\mathcal{B}_0$. ■

We will now apply the above material to the joint numerical range.

Theorem 1. Let $A_1, \dots, A_m \in \mathcal{M}_N$ be traceless, linearly independent hermitian operators. Let $\mathcal{A} \subset \mathcal{M}_N$ be the subspace spanned by them. Then i) The joint numerical range of $A_1, \dots, A_m$ is affinely isomorphic to $Pr_{\mathcal{A}}(\Omega_N)$; ii) The convex hull of the joint numerical range of $A_1, \dots, A_m$ is affinely isomorphic to $Pr_{\mathcal{A}}(\mathcal{Q}_N)$.

Proof. Let $U = \mathcal{M}_N, V = \mathbb{R}^m$. Let $v_1, \dots, v_m$ be the standard basis in $\mathbb{R}^m$. Then the map $jnr_{A_1, \dots, A_m}$ has the form equation (4) with $u_i = A_i$. By Proposition 1, the map $M : \mathcal{A} \rightarrow \mathbb{R}^m$ is a linear isomorphism. Set $G = \{X \in \mathcal{M}_N : \text{tr}(X) = 1\}$. Then $G \subset \mathcal{M}_N$ is an affine hyperplane containing the affine subspace $\mathcal{A} + \mathbf{1}_N$. Claim i) now follows from Corollary 1.

The convex hulls of affinely isomorphic sets are affinely isomorphic. Hence, claim i) implies claim ii). ■

Corollary 2. *Let the setting be as in Theorem 1, with $m = N^2 - 1$. Then $W(A_1, \dots, A_m)$ is affinely isomorphic to Ω_N and $\text{conv}[W(A_1, \dots, A_m)]$ is affinely isomorphic to $\mathcal{Q}_N$.*

Proof. *This is a special case of Theorem 1. In this case the affine hyperplane G is parallel to $\mathcal{A}$. Hence $\text{Pr}_{\mathcal{A}}$ induces an affine isomorphism of Ω_N and $\text{Pr}_{\mathcal{A}}(\Omega_N)$. ■*

Example 1. Let $N = 2$. The Pauli matrices

$$\sigma_1 = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \quad \sigma_2 = \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix}, \quad \sigma_3 = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \quad (7)$$

form an orthogonal basis in the space of traceless Hermitian 2×2 matrices. By Corollary 2,¹ $W(\sigma_1, \sigma_2, \sigma_3) \subset \mathbb{R}^3$ is a sphere. Let ρ be a 2×2 density matrix. Its *Bloch vector representation* is given by

$$\rho = \frac{1}{2} \mathbf{1}_2 + \sum_{j=1}^3 \tau_j \sigma_j. \quad (8)$$

We have $\tau_j = \text{tr}(\sigma_j \rho)$; the *Bloch vector* $(\tau_1, \tau_2, \tau_3) = \tau \in \mathbf{R}^3$ determines ρ . The positivity condition $\rho \geq 0$, implies

$$\sqrt{\tau_1^2 + \tau_2^2 + \tau_3^2} = \|\tau\| \leq 1/2.$$

The equality $\|\tau\| = 1/2$ holds iff $\rho = \rho_\psi$, a pure state. Thus, the map $\text{jnr}_{(\sigma_1, \sigma_2, \sigma_3)}$ yields an isomorphism of $\mathcal{Q}_2$ (resp. Ω_2) and the *Bloch ball* (resp. Bloch sphere).

We will now discuss a generalization of Example 1 to $N > 2$. Note that $d = N^2 - 1$ is the dimension of the space of traceless $N \times N$ hermitian matrices endowed with the scalar product equation (2). Let $\lambda_1, \dots, \lambda_d$ be a basis in this space. It is convenient to use an orthogonal, but not necessarily an orthonormal basis. For example, for $N = 3$ it is standard to use the Gell-Mann matrices $(\lambda_1, \dots, \lambda_8)$. They satisfy the orthogonality relations $\text{tr}(\lambda_j \lambda_k) = 2\delta_{jk}$ [19]. Let ρ be a $N \times N$ state. The counterpart of equation (8) is

$$\rho = \frac{1}{N} \mathbf{1}_N + \sum_{j=1}^d \tau_j \lambda_j. \quad (9)$$

The generalized Bloch vector $\tau = \tau(\rho)$ has d components $\tau_j = \text{Tr}(\rho \lambda_j) / \text{Tr}(\lambda_j^2)$. Assume now that the basis $\lambda_1, \dots, \lambda_d$ is orthonormal. Then the generalized Bloch vector τ_ψ of a pure state $\rho_\psi = |\psi\rangle\langle\psi|$ satisfies $\tau_j = \langle\psi|\lambda_j|\psi\rangle$. Hence the joint numerical range $W(\lambda_1, \dots, \lambda_d)$ contains the Bloch vector τ_ψ .

By Corollary 2, the convex hull $\text{conv}[W(\lambda_1, \dots, \lambda_d)]$ is affinely isomorphic to the set $\mathcal{Q}_N$ of quantum states. By Theorem 1, if $\lambda_1, \dots, \lambda_m$ are linearly independent, then the joint numerical range $W(\lambda_1, \dots, \lambda_m)$ is affinely isomorphic to a projection of $\mathcal{Q}_N$ to $\mathbf{R}^m$. This is a generalization of a result in [12].

3. Quantum maps

We recall that the set $\mathcal{L}_N$ of operators on $\mathcal{H}_N$ is a Hilbert space with the scalar product $(A, B) = \text{tr}(AB^*)$. Let $\mathcal{LL}_N$ be the space of linear operators on $\mathcal{L}_N$. For

¹See also [17, 12].

$\Phi \in \mathcal{LL}_N$ we denote by $\Phi^* \in \mathcal{LL}_N$ its adjoint. For $X, Y \in \mathcal{L}_N$ we define $\Phi_{X|Y} \in \mathcal{LL}_N$ by

$$\Phi_{X|Y}(A) = XAY^*. \quad (10)$$

The following lemma will be used in the proof of Proposition 3. The proof is straightforward and we leave it to the reader.

Lemma 2. 1. *The operators $\Phi_{X|Y}$ span the vector space $\mathcal{LL}_N$.*
 2. *Let $X_1, Y_1, X_2, Y_2 \in \mathcal{L}_N$. Then*

$$\Phi_{X_2|Y_2} \Phi_{X_1|Y_1} = \Phi_{X_2 X_1 | Y_2 Y_1}.$$

3. *We have*

$$(\Phi_{X|Y})^* = \Phi_{X^*|Y^*}.$$

We will denote by $\mathbf{1} \in \mathcal{L}_N$ the identity operator.

Definition 2. Let $k \geq 1$. We say that the k -tuple of operators $X_1, \dots, X_k \in \mathcal{L}_N$ is an identity resolution if

$$\sum_{i=1}^k X_i X_i^* = \mathbf{1}. \quad (11)$$

The dual property

$$\sum_{i=1}^k X_i^* X_i = \mathbf{1}, \quad (12)$$

holds iff $X_1^*, \dots, X_k^*$ is an identity resolution. Let $X_1, \dots, X_k \in \mathcal{L}_N$ be arbitrary. Set

$$\Phi_{(X_1, \dots, X_k)} = \sum_{i=1}^k \Phi_{X_i | X_i}. \quad (13)$$

The following lemma summarizes the basic properties of operators $\Phi_{(X_1, \dots, X_k)}$.

Lemma 3. Let $X_1, \dots, X_k \in \mathcal{L}_N$ be arbitrary, and let $\Phi = \Phi_{(X_1, \dots, X_k)} \in \mathcal{LL}_N$. Then the following holds.

1. *The operator $\Phi : \mathcal{L}_N \rightarrow \mathcal{L}_N$ preserves the sets $\mathcal{M}_N$ and $\mathcal{P}_N$.*
2. *The operators $X_1, \dots, X_k$ satisfy equation (11) iff $\Phi : \mathcal{L}_N \rightarrow \mathcal{L}_N$ preserves $\mathbf{1}$.*
3. *The operators $X_1, \dots, X_k$ satisfy equation (12) iff $\Phi : \mathcal{L}_N \rightarrow \mathcal{L}_N$ preserves the trace iff Φ^* preserves $\mathbf{1}$.*
4. *If $X_1, \dots, X_k$ satisfy equation (12), then $\Phi : \mathcal{L}_N \rightarrow \mathcal{L}_N$ preserves $\mathcal{Q}_N$.*

Proof. It suffices to prove 1) for operators $\Phi = \Phi_{X|X}$. The former property is immediate from $(XAX^*)^* = XA^*X^*$, and the latter from $\langle XAX^*v, v \rangle = \langle AX^*v, X^*v \rangle$. Claim 2) is immediate from the definition of Φ . We have

$$\text{Tr} \Phi(A) = (\Phi(A), \mathbf{1}) = (A, \Phi^*(\mathbf{1})).$$

Claim 3) now follows from Lemma 2. Claim 4) is immediate from 1) and 3). ■

For $\psi \in \mathcal{H}_N$ we set $\rho_\psi = |\psi\rangle\langle\psi|$. Then $\rho_\psi \in \mathcal{P}_N$, and for any $a \in \mathbb{C}$ we have

$$\rho_{a\psi} = |a|^2 \rho_\psi. \quad (14)$$

In particular, $\rho_\psi = 0$ iff $\psi = 0$ and $\rho_\psi \in \mathcal{Q}_N$ iff $\|\psi\| = 1$.

Proposition 2. Let $X_1, \dots, X_k \in \mathcal{L}_N$, and let $\Phi = \Phi_{(X_1, \dots, X_k)}$. Let $\psi \in \mathcal{H}_N$ be any vector. For $1 \leq i \leq k$ such that $X_i\psi \neq 0$, set $\psi_i = X_i\psi / \|X_i\psi\|$.

Then

$$\Phi(\rho_\psi) = \sum_i \|X_i\psi\|^2 \rho_{\psi_i}. \quad (15)$$

Proof. Let $X, Y \in \mathcal{L}_N$ and $v \in \mathcal{H}_N$. Then

$$(\Phi_{X|Y}(\rho_\psi))v = X\langle\psi, Y^*v\rangle\psi = \langle Y\psi, v\rangle X\psi.$$

In particular, we have

$$\Phi_{X|X}(\rho_\psi) = \rho_{X\psi}. \quad (16)$$

The claim now follows from equations (13) and (14). ■

Corollary 3. Let $X_1, \dots, X_k$ satisfy equation (12) and set $\Phi = \Phi_{(X_1, \dots, X_k)}$. Let $\psi \in \mathcal{H}_N$ be a unit vector. Let $1 \leq m \leq k$ be the number of unit vectors ψ_i from Proposition 2. Then

$$\Phi(\rho_\psi) = \sum_{i=1}^m p_i \rho_{\psi_i},$$

where $p_1, \dots, p_m > 0$ and $\sum p_i = 1$.

Proof. Set $p_i = \|X_i\psi\|^2$. The claim follows from equation (15) and the identity

$$\sum_i \|X_i\psi\|^2 = \left\langle \left(\sum_{i=1}^k X_i^* X_i \right) \psi, \psi \right\rangle.$$

■

Corollary 1. Let $X_1, \dots, X_k \in \mathcal{L}_N$, and set $\Phi = \Phi_{(X_1, \dots, X_k)}$. Suppose that $X_1, \dots, X_k$ satisfy equation (11). Then

1. For any $A \in \mathcal{L}_N$ we have

$$W(\Phi(A)) \subset W(A); \quad (17)$$

2. For any $A_1, \dots, A_m \in \mathcal{M}_N$ we have

$$W(\Phi(A_1), \dots, \Phi(A_m)) \subset \text{conv}(W(A_1), \dots, W(A_m)); \quad (18)$$

3. If $W(A_1, \dots, A_m)$ is convex, then we have

$$W(\Phi(A_1), \dots, \Phi(A_m)) \subset W(A_1, \dots, A_m). \quad (19)$$

Proof. Let $A \in \mathcal{L}_N$ be arbitrary, and let $\psi \in \mathcal{H}_N$ be a unit vector. Then

$$\mathrm{Tr}(\Phi(A)\rho_\psi) = \mathrm{Tr}\left(\left(\sum_{i=1}^k X_i A X_i^*\right)\rho_\psi\right) = \mathrm{Tr}\left(A\left(\sum_{i=1}^k X_i^* \rho_\psi X_i\right)\right).$$

Note that the operators $X_1^*, \dots, X_k^*$ satisfy equation (12). By Proposition 2 and Corollary 1, there are unit vectors $\psi_j \in \mathcal{H}_N$ and probabilities p_j such that

$$\mathrm{Tr}(\Phi(A)\rho_\psi) = \sum_j p_j \mathrm{Tr}(A\rho_{\psi_j}) = \sum_j p_j \langle \psi_j | A | \psi_j \rangle. \quad (20)$$

Claim 2 now follows from equation (3). Claim 3 is a special case of Claim 2. Claim 3 and the Toeplitz-Hausdorff theorem yield Claim 1. ■

Let $\Phi = \Phi_{(X_1, \dots, X_k)}$. In the physics literature the transformations of $\mathcal{P}_N$ defined by

$$\rho_1 = \Phi(\rho) = \sum_{i=1}^k X_i \rho X_i^* \quad (21)$$

are called *quantum maps*. Operators $X_i \in \mathcal{L}_N$ are the *measurement operators* or *Kraus operators*. Quantum maps correspond to generalized quantum measurements with k possible outcomes [18]. By Lemma 3, the map $\Phi : \rho \mapsto \rho_1$ is *unital* (resp. *trace preserving*) iff equation (11) (resp. equation (12)) is satisfied. The quantum map $\Phi^* = \Phi_{(X_1^*, \dots, X_k^*)}$ is dual to Φ . The duality of quantum maps corresponds to the duality between the Schrödinger and the Heisenberg representations in quantum mechanics. In the former, the quantum states $\rho \in \mathcal{P}_N$ evolve via $\rho_1 = \Phi(\rho)$, while the observables $A \in \mathcal{M}_N$ do not. In the latter, the quantum states do not change and the observables evolve by $A \mapsto A_1 = \Phi^*(A)$. The identity

$$\mathrm{Tr}(\Phi(\rho)A) = \mathrm{Tr}\left(\sum_i X_i \rho X_i^* A\right) = \mathrm{Tr}\left(\sum_i \rho X_i^* A X_i\right) = \mathrm{Tr}(\rho \Phi^*(A))$$

illustrates this duality. Note that Φ is trace preserving (resp. unital) iff Φ^* is unital (resp. trace preserving). Corollary 1 describes a relationship between quantum maps and numerical ranges. We will now present a few examples illustrating this relationship.

Example 2. Let $N = 2$. The *decaying channel* is the discrete dynamics $\Phi = \Phi_{X_1, X_2}$ corresponding to the Kraus operators $X_1 = \begin{bmatrix} 1 & 0 \\ 0 & \sqrt{1-p} \end{bmatrix}$ and $X_2 = \begin{bmatrix} 0 & \sqrt{p} \\ 0 & 0 \end{bmatrix}$, where $p \in (0, 1)$ is a free parameter. Set $A^{(j)} = \Phi^{j-1}(A^{(1)})$, where $A^{(1)} = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$. Note that $\mathrm{tr}(A^{(1)}) = 0$. The set $W(A^{(1)})$ is the disc of radius $1/2$ centered at the origin, i.e., at the *baricenter point* $\mathrm{tr}(A^{(1)})/2$. We have $A^{(j)} = (1-p)^{(j-1)/2} A^{(1)}$, and $W(A^{(j)}) = (1-p)^{(j-1)/2} W(A^{(1)})$ is the disc of radius $\frac{1}{2}(1-p)^{(j-1)/2}$ with the center at 0. For instance, $A^{(2)} = \Phi(A^{(1)}) = \begin{bmatrix} 0 & \sqrt{1-p} \\ 0 & 0 \end{bmatrix}$, and $A^{(3)} = \Phi^2(A^{(1)}) = \begin{bmatrix} 0 & 1-p \\ 0 & 0 \end{bmatrix}$. The




Figure 1: Numerical ranges $W(A^{(1)}), W(A^{(2)}), W(A^{(3)})$ from Example 2 for $p = 0.5$ and $W(B^{(1)}), W(B^{(2)}), W(B^{(3)})$ from Example 3 for $p = 0.25$. The stars mark the baricenter points in these examples. They are preserved by the dynamics.

limit of $W(A^{(j)})$, as $j \rightarrow \infty$, is $\{0\}$. Fig. 1a shows $W(A^{(1)}), W(A^{(2)}), W(A^{(3)})$ for $p = 0.5$.

Example 3. Let again $N = 2$. The *phase-flip channel* is the discrete dynamics $\Psi = \Phi_{X_1, X_2}$ corresponding to the Kraus operators $X_1 = \sqrt{1-p} \mathbf{1}_2$ and $X_2 = \sqrt{p} \sigma_1$. Here again, $p \in (0, 1)$ is a free parameter. We set $B^{(1)} = A^{(1)}$ and $B^{(j-1)} = \Psi^j(B^{(1)})$. Then $B^{(2)} = \begin{bmatrix} 0 & 1-p \\ p & 0 \end{bmatrix}$ and $B^{(3)} = \begin{bmatrix} 0 & 1-2p(1-p) \\ 2p(1-p) & 0 \end{bmatrix}$. The numerical ranges of all $B^{(j)}$ are ellipses. Fig. 1b shows $W(B^{(1)}), W(B^{(2)}), W(B^{(3)})$ for $p = 0.25$.

Example 4. This example is a generalization of Example 3 to $N = 3$. The *double flip channel acting on a qutrit* is the discrete dynamics $\Xi = \Phi_{X_1, X_2, X_3}$; the Kraus operators are $X_1 = \sqrt{1-p-q} \mathbf{1}_3$, $X_2 = \sqrt{p} \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ and $X_3 = \sqrt{q} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}$. The parameters p, q satisfy $0 \leq p, q, p+q \leq 1$. The operators X_2, X_3 correspond to bit flips with probabilities p and q . The operator Ξ is trace preserving. Let $C^{(1)} = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 2i \end{bmatrix}$. This

operator was studied in [16]; its numerical range $W(C^{(1)})$ is formed by the convex hull of an ellipse and a point outside it. The *baricenter* of $W(C^{(1)})$ is $\text{tr}(C^{(1)})/3 = (1+2i)/3$.

Set $C^{(j)} = \Xi^{j-1}(C^{(1)})$. For instance, $C^{(2)} = \begin{bmatrix} p & 1-p-q & q \\ p & 1-p-q(1-2i) & 0 \\ 0 & 0 & 2i+q(1-2i) \end{bmatrix}$.

In Figure 2 $p = 0.5$ and $q = 0.4$, hence $1-p-q = 0.1$ and all three Kraus operators contribute to the dynamics.




Figure 2: This figure illustrates Example 4 and the inclusion relation equation (17). It shows the numerical ranges $W(C^{(1)}) \supset W(C^{(2)}) \supset W(C^{(3)})$ for $p = 0.5, q = 0.4$. The limit of $W(C^{(j)})$, as $j \rightarrow \infty$, is the baricenter point $(1 + 2i)/3$ marked by the star in the figure.

4. Joint numerical shadows

The set $S(\mathcal{H}_N) = \{\psi \in \mathcal{H}_N : \|\psi\| = 1\}$ is the unit sphere S^{2N-1} . We denote by μ the normalized Haar measure on $S(\mathcal{H}_N)$, as well as its push-forward to $\Omega_N \simeq \mathbb{C}P^{N-1}$.² Let ν_A be the push-forward of μ to $\mathbb{C}$ under the numerical range map of an operator $A \in \mathcal{L}_N$. Then ν_A is the *numerical shadow* of A [14, 13]. Let dz denote the density of Lebesgue measure on $\mathbb{C}$. Then $d\nu_A(z) = P_A(z)dz$ where the probability distribution is given by

$$P_A(z) = \int_{\Omega_N} d\mu(\psi) \delta(z - \langle \psi | A | \psi \rangle). \quad (22)$$

Let now $A_1, \dots, A_m \in \mathcal{M}_N$, let $\text{jnr}_{(A_1, \dots, A_m)} : \Omega_N \rightarrow \mathbb{R}^m$ be the joint numerical range map, and let $W(A_1, \dots, A_m)$ be the JNR. The push-forward $\nu_{A_1, \dots, A_m}$ of μ to $\mathbb{R}^m$ under the map $\text{jnr}_{(A_1, \dots, A_m)}$ is the *joint numerical shadow* of $A_1, \dots, A_m$. Thus, $\nu_{A_1, \dots, A_m}$ is supported on $W(A_1, \dots, A_m)$. The corresponding probability distribution satisfies

$$P_{A_1, \dots, A_m}(x_1, \dots, x_m) = \int_{\Omega_N} d\mu(\psi) \prod_{j=1}^m \delta(x_j - \langle \psi | A_j | \psi \rangle). \quad (23)$$

Numerical shadow is the special case of the joint numerical shadow corresponding to $m = 2$. See the examples below for illustration.

If ν is a measure on $\mathbb{R}^m$ and $a \in \mathbb{R}$, we denote by ν^a the push-forward of ν under the self-map $v \mapsto av$ of $\mathbb{R}^m$. If ν_1, ν_2 are measures on $\mathbb{R}^m$, we denote by $\nu_1 * \nu_2$ their *convolution*. The following proposition exposes a few basic properties of joint numerical shadows. We leave the proof to the reader.

²It is known as the Fubini–Study measure in the physics literature.

Proposition 3. 1. Let $A_1, \dots, A_m \in \mathcal{M}_N$. Let $U \in \mathcal{L}_N$ be a unitary operator. Set $B_i = UA_iU^*$, $1 \leq i \leq m$. Then

$$\nu_{B_1, \dots, B_m} = \nu_{A_1, \dots, A_m}.$$

2. Let $A_1, \dots, A_m \in \mathcal{M}_N$ and $B_1, \dots, B_m \in \mathcal{M}_N$ be arbitrary. Let $a, b \in \mathbb{R}$. Then the joint numerical shadows satisfy

$$\nu_{aA_1+bB_1, \dots, aA_m+bB_m} = \nu_{A_1, \dots, A_m}^a * \nu_{B_1, \dots, B_m}^b.$$

Example 5. We review Example 1. The set $W(\sigma_1, \sigma_2, \sigma_3) \subset \mathbb{R}^3$ is the sphere of radius $1/2$. The joint numerical shadow $\nu_{\sigma_1, \sigma_2, \sigma_3}$ is the normalized Haar measure.

Example 6. We use the isomorphism $\mathcal{H}_4 = \mathcal{H}_2 \otimes \mathcal{H}_2$ to define the extensions of Pauli matrices $A_j = \sigma_j \otimes \mathbf{1}_2$, $j = 1, 2, 3$. We use the fact that the Haar measure on the space of pure states in $\mathcal{H}_A \otimes \mathcal{H}_B$ induces by partial trace, $\omega = \text{Tr}_B |\psi\rangle\langle\psi|$, the Lebesgue measure on the space of mixed states in $\mathcal{H}_2$ [20]. Using the equality between the expected values of an operator on $\mathcal{H}_2$ and the extended operator on $\mathcal{H}_4$, we obtain that ν_{A_1, A_2, A_3} is the normalized Lebesgue measure on the Bloch ball. Let $B_j = \mathbf{1}_2 \otimes \sigma_j$, $j = 1, 2, 3$. The *swap operator*

$$S = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}, \quad (24)$$

is a unitary operator on $\mathcal{H}_4$ satisfying $B_j = SA_jS^*$. By Proposition 3 and the above discussion, ν_{B_1, B_2, B_3} is the normalized Lebesgue measure on the Bloch ball.

Acknowledgements. This project started at the 43–th *Symposium on Mathematical Physics* in Toruń; it was accomplished during the 44–th Symposium. It is a pleasure to thank the organizers of these Symposia. We are obliged to Piotr Gawron for creating the figures. K. Ż. is grateful to C. F. Dunkl, J. A. Holbrook, J. Miszczał and Z. Puchała for fruitful discussions on numerical shadows and acknowledges financial support by the grant number N202 090239 of Polish Ministry of Science and Higher Education. The work of E.G. was partially supported by the MNiSzW grant N N201 384834. He acknowledges stimulating discussions with Bent Orsted in Aarhus in June 2011.

References

- [1] A. Horn and C. R. Johnson, *Topics in Matrix Analysis*, Cambridge University Press, Cambridge, 1994.
- [2] K. E. Gustafson and D. K. M. Rao, *Numerical Range: The Field of Values of Linear Operators and Matrices*. Springer-Verlag, New York, 1997.
- [3] O. Toeplitz, *Das algebraische Analogon zu einem Satze von Fejér*, Math. Zeitschrift **2**, 187–197 (1918).
- [4] F. Hausdorff, *Der Wertvorrat einer Bilinearform*, Math. Zeitschrift **3**, 314–316 (1919).
- [5] E. Gutkin, The Toeplitz-Hausdorff theorem revisited: relating linear algebra and geometry, *Math. Intelligencer* **26**, 8 – 14 (2004).
- [6] E. Jonckheere, F. Ahmad, E. Gutkin, Differential topology of numerical range, *Lin. Alg. Appl.* **279**, 227 – 254 (1998).

- [7] C. K. Li, A simple proof of the elliptical range theorem, *Proc. Am. Math. Soc.* **124**, 1985-1986 (1996).
- [8] D.S. Keeler, L. Rodman and I.M. Spitkovsky, The numerical range of 3×3 matrices, *Lin. Alg. Appl.* **252**, 115-1139 (1997).
- [9] D. W. Kribs, A. Pasieka, M. Laforest, C. Ryan, and M. P. Silva, Research problems on numerical ranges in quantum computing, *Linear and Multilinear Algebra* **57**, 491-502 (2009).
- [10] T. Schulte-Herbrüggen, G. Dirr, U. Helmke, and S. J. Glaser, The significance of the c -numerical range and the local c -numerical range in quantum control and quantum information, *Linear and Multilinear Algebra* **56**, 3-26 (2008)
- [11] P. Gawron, Z. Puchała, J. A. Miszczak, Ł. Skowronek, M.-D. Choi, and K. Życzkowski, Restricted numerical range: a versatile tool in the theory of quantum information, *J. Math. Phys.* **51**, 102204 (24pp) (2010).
- [12] C.F. Dunkl, P. Gawron, J.A. Holbrook, J. Miszczak, Z. Puchała and K. Życzkowski, Numerical shadow and geometry of quantum states, *J. Phys. A* **44**, 335301 (19pp) (2011).
- [13] T. Gallay and D. Serre, Numerical measure of a complex matrix, *Commun. Pure Appl. Math.* **65**, 287-336 (2012).
- [14] C.F. Dunkl, P. Gawron, J.A. Holbrook, Z. Puchała and K. Życzkowski, Numerical shadows: Measures and densities on the numerical range, *Lin. Algebra Appl* **434**, 2042-2080 (2011).
- [15] K. Życzkowski (with M.-D. Choi, C. Dunkl, J. Holbrook, P. Gawron, J. Miszczak, Z. Puchała, and Ł. Skowronek), Generalized numerical range as a versatile tool to study quantum entanglement, *Oberwolfach Report* **59**, 34-37 (2009).
- [16] Z. Puchała, J. A. Miszczak, P. Gawron, C. F. Dunkl, J. A. Holbrook, and K. Życzkowski, Restricted numerical shadow and geometry of quantum entanglement, *preprint* arXiv:1201.2524
- [17] E. Gutkin, E.A. Jonckheere, M. Karrow, Convexity of the joint numerical range: topological and differential geometric viewpoints, *Lin. Alg. Appl.* **376**, 143 – 171 (2004).
- [18] I. Bengtsson and K. Życzkowski, *Geometry of Quantum States*, Cambridge UP, Cambridge, 2006.
- [19] L. I. Schiff, *Quantum Mechanics*, New York, McGraw-Hill 1968.
- [20] K. Życzkowski and H.-J. Sommers, Induced measures in the space of mixed quantum states, *J. Phys. A* **34**, 7111-7125 (2001).